

Carsten Müssig eLiS UP

ENZYMES

Biochemistry I
Unit 7th

CONTENTS

- Chemistry
- Classification
- Mechanism of Enzyme Action
- Enzyme Kinetics
- Inhibition
- Activation
- Specificity






CHEMISTRY

Introduction

- Enzymes are *biological catalysts* that speed up the rate of the biochemical reaction.
- Most enzymes are three dimensional *globular proteins* (tertiary and quaternary structure).
- Some special RNA species also act as enzymes and are called *Ribozymes* e.g. hammerhead ribozyme.



Hammerhead enzyme


STRUCTURE OF ENZYMES

- The *active site* of an enzyme is the region that binds substrates, co-factors and prosthetic groups and contains residue that helps to hold the substrate.
- Active sites generally occupy less than 5% of the total surface area of enzyme.
- Active site has a *specific shape* due to tertiary structure of protein.
- A change in the shape of protein affects the shape of active site and function of the enzyme.




ACTIVE SITE

- Active site can be further divided into:



CO-FACTORS

- Co-factor is the non protein molecule which carries out chemical reactions that can not be performed by standard 20 amino acids.
- Co-factors are of two types:
 - Organic co-factors
 - Inorganic cofactors



INORGANIC CO-FACTORS

- These are the inorganic molecules required for the proper activity of enzymes.

Examples:

- Enzyme carbonic anhydrase requires Zn^{++} for its activity.
- Hexokinase has co-factor Mg^{++}

ORGANIC CO-FACTORS

- These are the organic molecules required for the proper activity of enzymes.

Example:


- Glycogen phosphorylase requires the small organic molecule pyridoxal phosphate.



TYPES OF ORGANIC CO-FACTORS


Prosthetic Group

- A prosthetic group is a permanently bound organic co-factor. E.g. NAD⁺ groups and biotin.




Coenzyme

- A coenzyme is loosely bound organic co-factor. E.g. Flavins, heme




- An enzyme with its co-factor removed is designated as *apoenzyme*.
- The complete complex of a protein with all necessary small organic molecules, metal ions and other components is termed as *holoenzyme* or *holoprotein*.




SUBSTRATE

- The reactant in biochemical reaction is termed as **substrate**.
- When a substrate binds to an enzyme it forms an **enzyme-substrate complex**.



SITES OF ENZYME SYNTHESIS

- Enzymes are synthesized by *ribosomes* which are attached to the rough endoplasmic reticulum.
- Information for the synthesis of enzyme is *carried by DNA*.
- Amino acids are bonded together to form specific enzyme according to the DNA's codes.



INTRACELLULAR AND EXTRACELLULAR ENZYMES

- **Intracellular** enzymes are *synthesized and retained in the cell for the use of cell itself.*
- They are found in the cytoplasm, nucleus, mitochondria and chloroplast.

Example :

- Oxydoreductase catalyses biological oxidation.
- Enzymes involved in reduction in the mitochondria.

- **Extracellular** enzymes are *synthesized in the cell but secreted from the cell to work externally.*

Example :

- Digestive enzyme produced by the pancreas, are not used by the cells in the pancreas but are transported to the duodenum.



CHARACTERISTICS

- Enzymes *speed up* the reaction by lowering the activation energy of the reaction.
- Their presence *does not effect* the nature and properties of *end product*.
- They are *highly specific* in their action that is each enzyme can catalyze one kind of substrate.
- Small amount of enzymes can accelerate chemical reactions.
- Enzymes are *sensitive* to change in pH, temperature and substrate concentration.
- *Turnover number* is defined as the number of substrate molecules transformed per minute by one enzyme molecule.

Catalase turnover number = 6×10^6 /min



NOMENCLATURE OF ENZYMES

- An enzyme is named according to the name of the substrate it catalyses.
- Some enzymes were named before a systematic way of naming enzyme was formed.

Example: pepsin, trypsin and rennin

- By adding suffix **-ase** at the end of the name of the substrate, enzymes are named.
- Enzyme for catalyzing the hydrolysis is termed as hydrolase.

Example :



EXAMPLES

substrate	enzymes	products
lactose	lact ase	glucose + galactose
maltose	malt ase	Glucose
cellulose	cellul ase	Glucose
lipid	lip ase	Glycerol + fatty acid
starch	amyl ase	Maltose
protein	prote ase	Peptides + polypeptide





CLASSIFICATION

CLASSIFICATION OF ENZYMES

- A systematic classification of enzymes has been developed by *International Enzyme Commission*.
- This classification is based on the type of reactions catalyzed by enzymes.
- There are *six* major classes.
- Each class is further divided into sub classes, sub sub-classes and so on, to describe the huge number of different enzyme-catalyzed reactions.



Classification of enzymes

Continued.....

ENZYME CLASS	REACTION TYPE	EXAMPLES
Oxidoreductases	Reduction-oxidation (redox)	Lactate dehydrogenase
Transferases	Move chemical group	Hexokinase
Hydrolases	Hydrolysis; bond cleavage with transfer of functional group of water	Lysozyme
Lysases	Non-hydrolytic bond cleavage	Fumarase
Isomerases	Intramolecular group transfer (isomerization)	Triose phosphate isomerase
Ligases	Synthesis of new covalent bond between substrates, using ATP hydrolysis	RNA polymerase





MECHANISM OF ENZYME ACTION

MECHANISM OF ENZYME ACTION

- The catalytic efficiency of enzymes is explained by two perspectives:


Thermodynamic
changes

Processes at the
active site



THERMODYNAMIC CHANGES

- All chemical reactions have energy barriers between reactants and products.
- The difference in transitional state and substrate is called *activation barrier*.




THERMODYNAMIC CHANGES


- Only a few substances cross the activation barrier and change into products.
- That is why rate of uncatalyzed reactions is much slow.
- Enzymes provide an alternate pathway for conversion of substrate into products.
- Enzymes accelerate reaction rates by forming transitional state having low activation energy.
- Hence, the reaction rate is increased many folds in the presence of enzymes.
- The total energy of the system remains the same and equilibrium state is not disturbed.



THERMO-DYNAMIC CHANGES OVERVIEW




PROCESSES AT THE ACTIVE SITE




COVALENT CATALYSIS

- Enzymes form covalent linkages with substrate forming transient enzyme-substrate complex with very low activation energy.
- Enzyme is released unaltered after completion of reaction.




ACID-BASE CATALYSIS

- Mostly undertaken by oxido- reductases enzyme.
- Mostly at the active site, histidine is present which act as both proton donor and proton acceptor.




CATALYSIS BY PROXIMITY

- In this catalysis molecules must come in bond forming distance.
- When enzyme binds:
 - A region of high substrate concentration is produced at active site.
 - This will orient substrate molecules especially in a position ideal for them.




CATALYSIS BY BOND STRAIN

- Mostly undertaken by *lyases*.
- The enzyme-substrate binding causes *reorientation* of the structure of site due to in a strain condition.
- Thus *transitional state* is required and here bond is unstable and eventually broken.
- In this way bond between substrate is *broken* and converted into *products*.




LOCK AND KEY MODEL

- Proposed by EMIL FISCHER in 1894.
- Lock and key hypothesis assumes the active site of an enzyme is rigid in its shape.
- There is no change in the active site before and after a chemical reaction.




INDUCED FIT MODEL

- More recent studies have revealed that the process is much more likely to involve an induced fit model(proposed by DANIAL KOSH LAND in 1958).
- According to this exposure of an enzyme to substrate cause a change in enzyme, which causes the active site to change it"s shape to allow enzyme and substrate to bind.



INDUCED FIT MODEL





ENZYMES KINETICS

INTRODUCTION

“It is a branch of biochemistry in which we study *the rate of enzyme catalyzed reactions.*”

- Kinetic analysis reveals the number and order of the individual steps by which enzymes transform substrate into products
- Studying an enzyme's kinetics in this way can reveal the catalytic mechanism of that enzyme, its role in metabolism, how its activity is controlled, and how a drug or an agonist might inhibit the enzyme



RATES OF REACTION AND THEIR DEPENDENCE ON ACTIVATION ENERGY

- Activation Energy (E_a):

“The least amount of energy needed for a chemical reaction to take place.”


- Enzyme (as a catalyst) acts on substrate in such a way that they lower the activation energy by changing the route of the reaction.
- The reduction of activation energy (E_a) increases the amount of reactant molecules that achieve a sufficient level of energy, so that they reach the activation energy and form the product.

Example:

- *Carbonic anhydrase* catalyses the hydration of 10^6 CO_2 molecules per second which is 10^7 x faster than spontaneous hydration.



ENZYMES LOWER THE ACTIVATION ENERGY OF A REACTION



KINETICS OF ENZYMES CATALYSIS

- Enzymes catalysis:

“ It is an increase in the rate of reaction with the help of enzyme(as catalyst).”

- Catalysis by enzymes that proceed via unique reaction mechanism, typically occurs when the transition state intermediate forms a covalent bond with the enzyme(covalent catalysis).
- During the process of catalysis enzymes always emerge unchanged at the completion of the reaction.



FACTORS AFFECTING RATE OF ENZYME CATALYZED REACTIONS

- Temperature
- Hydrogen ion concentration(pH)
- Substrate concentration



EFFECT OF TEMPERATURE


- Raising the temperature increases the rate of enzyme catalyzed reaction by increasing kinetic energy of reacting molecules.
- Enzymes work maximum over a particular temperature known as *optimum temperature*. Enzymes for humans generally exhibit stability temperature up to 35-45 ° C.
- The temperature coefficient is a factor Q_{10} by which the rate of biological processes increases for a 10 ° C increase in temperature.
- For most biological processes $Q_{10} = 2$.
- However some times heat energy can also increase kinetic energy to a point that exceed the energy barrier which results in denaturing of enzymes.



Temperature

5- 40°C

Increase in Activity




40°C - denatures

<5°C - inactive




EFFECT OF PH

- Rate of almost all enzymes catalyzed reactions depends on pH
- Most enzymes exhibit optimal activity at pH value between *5 and 9*
- High or low pH value than optimum value will cause ionization of enzyme which result in denaturation of enzyme



PH AFFECTS THE FORMATION OF HYDROGEN BONDS AND SULPHUR BRIDGES IN PROTEINS AND SO AFFECTS SHAPE.



MICHAELIS-MENTEN MODEL & EFFECTS OF SUBSTRATE CONCENTRATION

- Michaelis-Menten Model:

“According to this model the enzyme reversibly combines with substrate to form an ES complex that subsequently yields product, regenerating the free enzyme.”



where:

- S is the substrate
- E is the enzyme
- ES-is the enzyme substrate complex
- P is the product
- K1,K-1 and K2 are rate constants



MICHAELIS-MENTEN EQUATION

- Michaelis-Menten Equation:

“It is an equation which describes how reaction velocity varies with substrate concentration.”

$$V_o = \frac{V_{\max} [S]}{K_m + [S]}$$

- Where
 - V_o is the initial reaction velocity.
 - V_{\max} is the maximum velocity.
 - K_m is the Michaelis constant = $(k_{-1} + k_2)/k_1$.
 - $[S]$ is the substrate concentration.




ASSUMPTIONS FOR MICHAELIS-MENTEN EQUATION

- Following assumptions are made in deriving the Michaelis-Menten equation:
 - Relative concentrations of E and S.
 - Steady-State assumptions
 - Initial Velocity




SUBSTRATE CONCENTRATION



SUBSTRATE CONCENTRATION

Michaelis Menten Plot

$$V = \frac{V_{\max} \cdot [S]}{K_m + [S]}$$



PHARMACEUTICAL IMPORTANCE

- Enzymes are virtually involved in all physiological processes which makes them the *targets of choice for drugs* that cure or ameliorate human disease.
- Applied enzyme kinetics represents the *principal tool* by which scientist identify and characterize therapeutic agents that selectively inhibit the rates of specific enzymes catalyzed processes.
- Enzymes kinetics thus play a critical role in *drug discovery* as well as elaborating the mode of action of drugs.






INHIBITION

INHIBITION


- The prevention of an enzyme process as a result of interaction of inhibitors with the enzyme.

- **INHIBITORS:**

Any substance that can diminish the velocity of an enzyme catalyzed reaction is called an inhibitor.



TYPES OF INHIBITION



REVERSIBLE INHIBITION

- It is an inhibition of enzyme activity in which the inhibiting molecular entity can associate and dissociate from the protein's binding site.


TYPES OF REVERSIBLE INHIBITION

- There are four types:
 - Competitive inhibition.
 - Uncompetitive inhibition.
 - Mixed inhibition.
 - Non-competitive inhibition.



COMPETITIVE INHIBITION

- In this type of inhibition, the inhibitors compete with the substrate for the active site. Formation of E.S complex is reduced while a new E.I complex is formed.




EXAMPLES OF COMPETITIVE INHIBITION

- Statin Drug As Example Of Competitive Inhibition:
 - Statin drugs such as *lipitor* compete with HMG-CoA(substrate) and inhibit the active site of *HMG CoA-REDUCTASE* (that bring about the catalysis of cholesterol synthesis).



UNCOMPETITIVE INHIBITION

- In this type of inhibition, inhibitor does not compete with the substrate for the active site of enzyme instead it binds to another site known as *allosteric* site.



EXAMPLES OF UNCOMPETITIVE INHIBITION

- Drugs to treat cases of poisoning by methanol or ethylene glycol act as uncompetitive inhibitors.
- Tetramethylene sulfoxide and 3-butylthiolene 1-oxide are uncompetitive inhibitors of liver alcoholdehydrogenase.



MIXED INHIBITION

- In this type of inhibition both E.I and E.S.I complexes are formed.
- Both complexes are catalytically inactive.


NON COMPETITIVE INHIBITION

- It is a special case of inhibition.
- In this inhibitor has the same affinity for either enzyme E or the E.S complex.



IRREVERSIBLE INHIBITION

- This type of inhibition involves the *covalent attachment* of the inhibitor to the enzyme.
- The *catalytic activity* of enzyme is completely lost.
- It can only be restored only by synthesizing molecules.



EXAMPLES OF IRREVERSIBLE INHIBITION

- *Aspirin* which targets and covalently modifies a key enzyme involved in inflammation is an irreversible inhibitor.
- **SUICIDE INHIBITION :**
 - It is an unusual type of irreversible inhibition where the enzyme converts the inhibitor into a reactive form in its active site.





ACTIVATION

ACTIVATION

- Activation is defined as the conversion of an inactive form of an enzyme to active form which processes the metabolic activity.

TYPES OF ACTIVATION

- Activation by co-factors.
- Conversion of an enzyme precursor.



ACTIVATION BY CO FACTORS

- Many enzymes are activated by co-factors.

Examples:

- *DNA polymerase* is a holoenzyme that catalyzes the polymerization of de -oxyribonucleotide into a DNA strand. It uses Mg- ion for catalytic activity.
- *Horse liver dehydrogenase* uses Zn- ion for it"s activation.



CONVERSION OF AN ENZYME PRECURSOR


- Specific proteolysis is a common method of activating enzymes and other proteins in biological system.

Example:

- The generation of trypsin from trypsinogen leads to the activation of other zymogens.



ZYMOGEN ACTIVATION BY PROTEOLYTIC CLEAVAGE






ENZYME SPECIFICITY

ENZYME SPECIFICITY

- Enzymes are highly specific in nature, interacting with one or few substrates and catalyzing only one type of chemical reaction.
- Substrate specificity is due to complete fitting of active site and substrate .

Example:

- Oxydoreductase do not catalyze hydrolase reactions and hydrolase do not catalyze reaction involving oxidation and reduction.



TYPES OF ENZYME SPECIFICITY

- Enzymes show different degrees of specificity:
 - Bond specificity.
 - Group specificity.
 - Absolute specificity.
 - Optical or stereo-specificity.
 - Dual specificity.




BOND SPECIFICITY

- In this type, enzyme acts on substrates that are similar in structure and contain the same type of bond.

Example :

- **Amylase** which acts on α -1-4 glycosidic bond in starch dextrin and glycogen, shows bond specificity.




GROUP SPECIFICITY

- In this type of specificity, the enzyme is specific not only to the type of bond but also to the structure surrounding it.

Example:

- *Pepsin* is an endopeptidase enzyme, that hydrolyzes central peptide bonds in which the amino group belongs to aromatic amino acids e. g phenyl alanine, tyrosine and tryptophan.




SUBSTRATE SPECIFICITY


- In this type of specificity ,the enzymes acts only on one substrate

Example :

- *Uricase* ,which acts only on uric acid, shows substrate specificity.



- *Maltase* , which acts only on maltose, shows substrate specificity.




OPTICAL / STEREO-SPECIFICITY


- In this type of specificity, the enzyme is not specific to substrate but also to its optical configuration

Example:

- D amino acid oxidase acts only on D amino acids.



- L amino acid oxidase acts only on L amino acids.




DUAL SPECIFICITY

- There are two types of dual specificity.
- The enzyme may act on one substrate by two different reaction types.

Example:

- *Isocitrate dehydrogenase* enzyme acts on isocitrate (one substrate) by oxidation followed by decarboxylation (two different reaction types) .



DUAL SPECIFICITY

- The enzyme may act on two substrates by one reaction type

Example:

- *Xanthine* oxidase enzyme acts on xanthine and hypoxanthine (two substrates) by oxidation (one reaction type)

